arXiv:1209.1834v1 [physics.flu-dyn] 9 Sep 2012

Swarms of particles settling under gravity in a viscous fluid

Maria L. Ekiel-Jeżewska
Institute of Fundamental Technological Research,
Polish Academy of Sciences, Pawińskiego 5B, 02-106 Warsaw, Poland
(Dated: September 11, 2012)

We investigate swarms made of a small number of particles settling under gravity in a viscous fluid. The particles do not touch each other and can move relative to each other. The dynamics is analyzed in the point-particle approximation. A family of swarms is found with periodic oscillations of all the settling particles. In the presence of an additional particle above the swarm, the trajectories are horizontally repelled from the symmetry axis, and flattened vertically. The results are used to explain how a spherical cloud, made of a large number of particles distributed at random, evolves and destabilizes.

I. INTRODUCTION

A large number of close but non-touching particles, randomly distributed in a spherical volume of a viscous fluid and settling under gravity in the same fluid at a low-Reynolds-number, forms a swarm, which evolves in a remarkable way. The particles circulate and stay together for a long time, with the average particle and fluid velocity fields the same as in case of a more dense fluid drop[1–4]. An initially spherical swarm slowly flattens and expands, forming a torus and occasionally leaving single particles behind (which, as slower, form a thin long 'tail' above the swarm). Then, suddenly, the swarm breaks into two (or sometimes more) fragments which repeat the same evolution pattern. Dynamics of such a system has been extensively investigated experimentally and numerically, with the use of different methods[3,5–13]. However, it is still not clear what is the reason for the observed evolution pattern.

The goal of this work is to construct and analyze simple models, which help to understand basic features of a swarm evolution. Sec. II contains the guiding questions and presentation of the theoretical framework. In Sec. III, a new family of periodic solutions is found and analyzed, as a model of the swarm. Sec. IV illustrates how these periodic motions are influenced by the presence of an additional particle (to model the tail influence on the swarm dynamics). Discussion of the results and conclusions are presented in Sec. V.

II. PROBLEM

We aim towards understanding the evolution pattern of a swarm of heavy particles settling in a viscous fluid at the Reynolds number much smaller than unity. We are going to investigate why a swarm slowly deforms (expands, flattens, forms a torus) and then suddenly breaks, by relating these processes to the circulation of particles inside the swarm, and to the occasional loss of the particles into a tail above the swarm. To this goal, we construct and analyze models which contain the characteristic features of the swarm dynamics (particle circulations, their interaction with a tail, deformation of the swarm shape and its destabilization), and find out how these properties are interrelated.

The theoretical background is the following. Assume that K point-particles, located at $\boldsymbol{r}_i(t)$, move in a fluid of viscosity η under identical gravitational forces $\boldsymbol{F}$. The fluid velocity $\mathbf{v}$ and pressure p satisfy the Stokes equations,

$$\eta \boldsymbol{\nabla}^2 \mathbf{v}(\boldsymbol{r}) - \boldsymbol{\nabla} p(\boldsymbol{r}) = -\boldsymbol{F} \sum_{i=1}^{K} \delta(\boldsymbol{r} - \boldsymbol{r}_i), \qquad \boldsymbol{\nabla} \cdot \mathbf{v}(\boldsymbol{r}) = 0.$$

In the reference frame moving with a single particle,

$$\dot{\boldsymbol{r}}_i(t) = \left[\sum_{k \neq i}^{K} \boldsymbol{\mu}_{ik} \right] \cdot \boldsymbol{F}, \qquad i = 1, ..., K, \tag{1}$$

where the mobility $\boldsymbol{\mu}_{ik}$ is given by the Oseen tensor[14],

$$\boldsymbol{\mu}_{ik} = \frac{1}{8\pi\eta r_{ik}} (\mathbf{I} + \hat{\boldsymbol{r}}_{ik}\hat{\boldsymbol{r}}_{ik}) \quad \text{for } i \neq k, \tag{2}$$

with $\hat{\boldsymbol{r}}_{ik} = (\boldsymbol{r}_i - \boldsymbol{r}_k)/r_{ik}$ and $r_{ik} = |\boldsymbol{r}_i - \boldsymbol{r}_k|$.

III. FAMILY OF PERIODIC SOLUTIONS

Hocking analyzed oscillations of four point-particles settling under gravity in a vertical plane[15]. His initial configurations can be modified to start from point particles located at vertices of a rectangle with vertical and horizontal sides, all located in the same vertical plane.

A straightforward 3D generalization results in initial configurations of $2N$ point-particles located at vertices of a regular right prism, which consists of 'twin' horizontal regular N-polygons, with one particle exactly above another one (an example is shown in Fig. 1). Such configurations lead to periodic relative motion of the particles in a vertical plane which includes the center-of-mass of the whole group, as schematically indicated in Fig. 1. In the reference frame of the center-of-mass, a particle and its twin follow the same periodic trajectory.

The length unit D is twice the initial distance of a particle from the symmetry axis, and the time unit is $8\pi\eta D^2/F$.



FIG. 1: Initial configuration of 16 particles (dots). A. Side view. B. Top view. Solid lines: planes of periodic motion.

As an example, we consider evolution of the initial configuration of 16 particles, shown in Fig. 1, with $c = 0.8$. The frame of reference is chosen in such a way that the z-axis is anti-parallel to gravity, and the xz-plane contains two periodic trajectories of 2 particles each. All the particle trajectories, observed in the laboratory frame of reference during $t = 25$, are plotted in Fig. 2.

There appear two characteristic stages of the evolution. In the first one, for $t \lesssim 3$ (five periods), the particles perform periodic oscillations while settling down. Then, the cluster destabilizes, and the motion is not regular any more. However, at least half of the particles stay close to each other until $t = 25$, and interact hydrodynamically, what is seen as curved fluctuating trajectories. Gradually, individual particles separate form the cluster, and as slower, are left behind. The second stage of the evolution is much longer than the first one.



FIG. 2: Trajectories of 16 particles initially located as shown in Fig. 1, with $c = 0.8$ (side view).

In the first stage of the evolution, owing to the rotational symmetry of the system, all the trajectories are planar, and their shapes are identical.

We now adopt the center-of-mass frame, and investigate how the shape of the periodic trajectory changes with the decrease of c. The trajectory height (just equal to c), width w, and its distance to the symmetry axis $h/2$, indicated in Fig. 3, are used as characteristic parameters



FIG. 3: Trajectories in the center-of-mass frame for $c{=}0.1$.

of the trajectories, together with the aspect ratio,

$$p = c/w. \tag{3}$$

In Fig. 4 we illustrate that the width w and the aspect ratio p of trajectories are getting smaller when c decreases down to $c{=}0.2$. Their length L decreases, and



FIG. 4: Trajectories in the center-of-mass frame, for large values of $c = 2.2, 1.4, 0.9, 0.6, 0.2$. The smaller c, the shorter the trajectory and the smaller its width.



FIG. 5: Trajectories in the center-of-mass frame, for small values of $c = 0.1, 0.05, 0.047$. The smaller c, the longer and wider the trajectory. Note a different scale on each axis.

a successive trajectory is located inside the previous one. When c decreases, the distance $h/2$ from a trajectory to the symmetry axis of the cluster ("radius of the hole") increases up to one - the upper limit determined by the initial conditions.



FIG. 6: The trajectory width, w, and twice its distance from the symmetry axis, h, versus c, for $c \geq 0.04481$. Inset: for $c \geq 0.8$, the ratio h/w increases linearly with $1/p$.

For small values $c < 0.2$, trajectories change their shape significantly. The length L and width w of trajectories increase rapidly when c is decreased by a tiny fraction, while h remains close to unity, see Figs. 5 and 6.

As shown in Fig. 7, the aspect ratio p of a trajectory (what can be interpreted as the aspect ratio of a cluster) decreases when c is decreased, and for $c \gtrsim 0.3$, the slope is smaller than for $c \lesssim 0.2$. The inset indicates that the trajectory width w becomes very large close to $c = c_0$.

Indeed, we observe that the length L and width w of a center-of-mass trajectory, and also period T of the oscillations increase when c decreases. Is there a critical value $c = c_0$, where they all become infinite? To check, we assume a power law divergence,

$$T \sim A/(c - c_0)^{\alpha} \text{ and } w \sim B/(c - c_0)^{\beta}, \quad \text{for } c \to c_0, \tag{4}$$

and search for c_0, α, β by plotting T and w versus $(c - c_0)$ in the log-log scale, see Fig. 8.



FIG. 7: Aspect ratio p of the cluster as a function of c (symbols). Solid straight lines: for larger c (red online), $p = 0.44c + 0.09$ and for small c (blue online), $p = c - c_0$.



FIG. 8: The period T and the cluster width w versus $c - c_0$. Solid lines: $T = 0.0013/(c - c_0)^{1.5}$ and $w = c_0/(c - c_0)$.

In this way, we find the critical value of c,

$$c_0 = 0.044788..., \tag{5}$$

and the critical exponents,

$$\alpha \approx 1.49 - 1.50, \qquad \beta \approx 1.00. \qquad (6)$$

For $c < c_0$, there is no periodic motions; the swarm form the beginning splits into N pairs of particles. The comparison of the trajectories without and with periodic oscillations is performed in Fig. 9. For $c < c_0$, the slope of a trajectory monotonically decreases to a constant value, which is smaller than the minimal slope reached at periodic trajectories.



FIG. 9: Two types of the particle trajectories: periodic oscillations ($c = 0.05, 0.049, 0.045$) and separation into pairs without oscillations ($c = 0.04, 0.03, 0.02$).

IV. A DROP AND A TAIL

We will now illustrate how periodic motions, investigated in the previous section, are modified in the presence of an additional particle. Such a system is supposed to model a suspension drop and a tail of particles gradually lost from it. It is known that the particles, which separate out from a suspension drop, circulate along the exterior trajectories up to the top of the drop, and escape along the symmetry axis[3].

In our model, the drop consists of 16 particles, and the tail just from a singlet. The initial configuration of 16 particles, located at vertices of a regular prism, is shown in Fig. 1, and we take $c = 0.9$. At $t = 0$, the 17th particle is placed at the symmetry axis of the prism, at a small distance $z_0 = 1.25$ above its center-of-mass. The question is how the trajectories of 16 particles, evaluated in their center-of-mass frame (and shown as the middle curves in Fig. 4), are modified by the 17$^{\text{th}}$ particle. Can a small tail change motion of a large drop?

The trajectories of the particles from our "drop" are shown in Fig. 10. The particles move in the plane yz of the plot. In the description of the axes, we indicated that the reference frame is moving with the center-of-mass of the 16 particles which form the "drop". It is clear that with time, the shape of the trajectory changes. It becomes less high, wider, and its distance from the symmetry axis increases. The same features are observed when the long-time part of the trajectory from Fig. 10 is compared to the corresponding trajectory in the absence of the 17$^{\text{th}}$ particle, see the curve with $c = 0.9$ in Fig. 4.



FIG. 10: Trajectories in the center-of-mass frame of 16 particles, which initially form a regular prism ($c = 0.9$), with an additional single particle on its symmetry axis as indicated.

The explanation is that the tail interacts hydrodynamically with the drop. The additional velocity of a drop particle, gained owing to its interaction with the tail (the Oseen velocity generated by the single point-force), is schematically indicated by arrows in Fig. 11. As the result, the particles inside the drop are repelled horizontally from the drop center, and attracted vertically to the central plane. The closer the tail, the larger the effect.



FIG. 11: Schematic explanation of the horizontal expansion and vertical flattening of the drop trajectories. Arrows (sketch): this part of the drop-particle velocity, which comes from its interaction with the tail.

Similar shape deformation of the trajectories is observed for spheres rather than points. The results for a drop and a tail made of 8+1 spherical particles are discussed in Appendix A.

V. DISCUSSION AND CONCLUSIONS

A new class of unstable periodic relative motions has been found: initially, $2N$ point-particles form a regular prism (two mirror horizontal regular polygons) with a vertical-to-horizontal aspect ratio c larger than a critical value c_0. (For $c < c_0$, the system from the beginning separates into pairs of the mirror particles.) Periodic motions have been observed for different values of N, with a smaller c_0 for a larger N; for example, $c_0 = 0.017$ for $2N = 32$. For larger values of N, it is difficult to investigate periodic motions, because they destabilize faster. For example, the swarms with $w \approx c$ and $N = 16, 32, 64$, start to move non-periodically after $\frac{5}{2}, \frac{5}{4}, \frac{1}{2}$ of the corresponding period ($T = 1.31, 0.52, 0.22$), respectively.

The mean settling velocity V_s of a swarm with $w \approx c$, averaged over its period, is close to the mean velocity $V_d = 2NF/(5\pi\eta w)$ of a fluid drop of the same weight and size, with the ratio $V_s/V_d = 0.81, 0.88, 0.97$ for the number of particles in the swarm $N = 16, 32, 64$, respectively.

It is supposed that the particle trajectories in the suspension drop are close to 'a periodic solution'. Particles are left behind the drop (as 'a tail') if sufficiently separated from these periodic orbits. Such a mechanism was confirmed in case of a swarm consisting of 3 particles (both point-like and spherical)[16–18]. It was shown that 3 close particles circulate together as a swarm for a long time before destabilizing into a faster pair and a slower singlet, and the interaction time is very sensitive to the initial conditions[16,17]. The observed chaotic scattering was associated with the existence of unstable periodic motions which for point-particles have not been found explicitly[16]. For 3 spherical particles with a very limited class of the initial conditions, such periodic trajectories of the relative motion were found[18]. For other initial conditions, the shape of the relative trajectories of the swarm particles indeed resembles the shape of the periodic ones[18].

It is expected that the pattern observed for 3 particles is generic also for many-particle systems. Periodic motions 'hidden' inside a suspension drop can correspond to periodic motions of a regular array made of a comparable number of particles, or a smaller number of effective particles. The regular swarm analyzed in Sec. III is just an example.

In Fig. 2, we have shown that a regular swarm does not break immediately after destabilization. On the contrary, most of its particles stay close to each other, and perform irregular circulations for a very long time. In this second stage of its evolution, the swarm clearly 'stays close' to the periodic regular solution from the first stage of the evolution.

Owing to statistical fluctuations, particles which are too far from periodic orbits, stop circulating and are left behind the drop as a tail above the symmetry axis of the drop. This mechanism is consistent with a wide range of the drop destabilization times, observed in experiments and simulations[13], just as it is in case of 3 particles only[16].

As demonstrated in Sec. IV for a simple model, the tail repels particles horizontally from the drop center and attracts them vertically to the central horizontal plane of the drop. This indicates that larger times t of the drop evolution correspond to smaller values of c for the model periodic solution of a regular swarm, analyzed in Sec. III. There is an analogy between the dynamics of the drop particles at an increasing time t and the dynamics of the regular swarm with the decreasing value of the parameter c. This analogy is restricted to a limited range of the c values, starting from $c \approx w$ (e.g. $c = 2.06$ for $N = 16$), and decreasing its value only a little, to match the drop aspect ratio $p \gtrsim 0.5 - 0.6$ observed in simulations and experiments[8]. For our model, this requirement excludes values of c comparable to the critical value c_0, see Fig. 7.

Destabilization time of a suspension drop can be associated with the characteristic destabilization time τ_d of the corresponding periodic orbits. The characteristic time scale τ_t of the drop deformation caused by the particle loss seems to be much larger than the characteristic destabilization time τ_d of the periodic solution.

In future publications, stability and other basic features of the periodic solutions presented here will be analyzed in details, using the methods applied elsewhere[19,20], and related to the corresponding properties of the point-particle simulations for suspension drops. Also, it is interesting to perform a comparison with periodic orbits for a regular system of settling non-spherical particles in a similar geometry, which have been recently found in simulations and experiments[21]. From both fundamental and practical point of view, it is important to confirm what is the physical mechanism of the drop destabilization.

Acknowledgments

This work was supported in part by the Polish Ministry of Science under grant 2011/01/B/ST3/05691.

Appendix A: Spherical particles

Consider now K identical spheres moving in a viscous fluid under gravitational forces $\boldsymbol{F}$. The fluid velocity $\mathbf{v}$ and pressure p satisfy the Stokes equations with the stick boundary conditions at the sphere surfaces. Dynamics of the translational motion of the spheres reads

$$\dot{\boldsymbol{r}}_i(t) \;=\; \left[\sum_{k=1}^{K} \boldsymbol{\mu}_{ik}\right] \cdot \boldsymbol{F}, \qquad i = 1, ..., N, \qquad \text{(A1)}$$

where $\boldsymbol{r}_i(t)$ are time-dependent positions of the sphere centers and the mobility matrices $\boldsymbol{\mu}_{ik}$ (which depend on relative positions of all the particles) are evaluated numerically by the multipole expansion[22,23] with the use of the HYDROMULTIPOLE numerical code[24].

To illustrate periodic gravitational settling of $2N$ identical spheres, we consider the initial configuration of their centers at vertices of a cube, with the side equal $1.5d$, where d is the sphere diameter (see the upper panel of Fig. 12). Differently from Fig. 4, we show projections of the trajectories on the plane of a cube wall (rather than the trajectories themselves). Circulations of the spheres in their center-of-mass frame are shown in the bottom panel of Fig. 12.

Similarly as in Sec. IV, we now investigate "a suspension drop with a tail", and we check how does the shape of the trajectories in Fig. 12 (drop) change in the presence of an additional single sphere (tail), initially located above the cube at its symmetry axis, as shown in the upper panel of Fig. 13.



FIG. 12: Model of a drop. Top: initial configuration of 8 spheres at vertices of a cube. Bottom: Trajectories of these sphere centers in their center-of-mass frame. (Projection on a vertical plane of a cube wall.)

In the bottom panel of Fig. 13, we show trajectories of the drop sphere centers in their center-of-mass frame after a long time, when the single sphere has been already left up far away. The comparison with Fig 12 shows that, owing to the hydrodynamic interaction with the tail, the width of the drop has become larger, the height smaller, and the hole radius has increased, in agreement with our findings in Sec. IV for the analogical point-particle system.

Evolution of the system in the reference frame of the drop is shown in the movie linked online to Fig. 13. It looks qualitatively the same as for the analogical system of point-particles studied in Sec. IV. Eight spheres circulate inside the drop, while the single tail sphere separates from the drop by a rapidly increasing distance.



FIG. 13: Model of a drop and a tail. Top: initial configuration of 8 spheres (drop) and one above it (tail). Bottom: Trajectories of the drop sphere centers in their center-of-mass frame after a time, during which the tail was left up far away. (Enhanced online).

---

[1] J. S. Hadamard. Mouvement permanent lent d'une sphere liquide et visqueuse dans un liquide visqueux. *C. R. Acad. Sci. (Paris)* **152**, 1735–1738 (1911).

[2] W. Rybczyński. Über die fortschreitende Bewegung einer flüssigen Kugel in einem zähen Medium. *Bull. Acad. Sci. Cracovie* **A**, 40–46 (1911).

[3] J. M. Nitsche and G. K. Batchelor. Break-up of a falling drop containing dispersed particles. *J. Fluid Mech.*, **340**, 161–175 (1997).

[4] M. L. Ekiel-Jeżewska, B. Metzger, and É. Guazzelli. Spherical cloud of point particles falling in a viscous fluid. *Phys. Fluids*, **18**, 038104 (2006).

[5] K. Adachi, S. Kiriyama, and N. Yoshioka. The behavior of a swarm of particles moving in a viscous fluid. *Chem. Eng.*

Sci. **33**, 115-121 (1978).

[6] G. Machu, W. Meile, L. C. Nitsche, and U. Schaflinger. Coalescence, torus formation and break-up of sedimenting drops: experiments and computer simulations. *J. Fluid Mech.*, **447**, 299–336 (2001).

[7] T. Bosse, L. Kleiser, C. Haertel, and E. Meiburg. Numerical simulation of finite Reynolds number suspension drops settling under gravity. *Phys. Fluids*, **17**, 037101 (2005).

[8] B. Metzger, M. Nicolas, and É. Guazzelli. Falling clouds of particles in viscous fluids. *J. Fluid Mech.*, **580**, 283–301 (2007).

[9] G. C. Abade and F. R. Cunha. Computer simulation of particle aggregates during sedimentation. *Comput. Methods Appl. Mech. Engrg.*, **196**, 4597–4612 (2007).

[10] G. Subramanian and D. L. Koch. Evolution of clusters of sedimenting low-Reynolds-number particles with Oseen interactions. *J. Fluid Mech.*, **603**, 63–100 (2008).

[11] S. Alabrudziński, M. L. Ekiel-Jeżewska, D. Chehata-Gòmez, and T. A. Kowalewski. Particle clusters settling under gravity in a viscous fluid. *Phys. Fluids*, **21** 073302 (2009).

[12] A. Myłyk, M. L. Ekiel-Jeżewska, How walls influence destabilization of a suspension drop settling under gravity in a viscous fluid?. Colloids Surf. A **365**, 109-111 (2010).

[13] A. Myłyk, W. Meile, G. Brenn, and M. L. Ekiel-Jeżewska, Break-up of suspension drops settling under gravity in a viscous fluid close to a vertical wall. Phys. Fluids **23**, 063302 (2011).

[14] S. Kim and S. J. Karrila. *Microhydrodynamics. Principles and Selected Applications.* Butterworth-Heinemann, London, 1991.

[15] L. M. Hocking, The behaviour of clusters of spheres falling in a viscous fluid. *J. Fluid Mech.* **20**, 129–139 (1964).

[16] I. M. Jànosi, T. Tèl, D. E. Wolf, and J. A. C. Gallas. Chaotic particle dynamics in viscous flows: The three-particle Stokeslet problem. *Phys. Rev. E*, **56**, 2858–2868 (1997).

[17] M. L. Ekiel-Jeżewska and E. Wajnryb, Lifetime of a cluster of spheres settling under gravity in Stokes flow, *Phys. Rev. E* **83**, 067301 (2011).

[18] M. L. Ekiel-Jeżewska, Periodic orbits of Stokesian dynamics, CDROM Proceedings of the XXII International Congress of Theoretical and Applied Mechanics, edited by J. Denier, M. D. Finn and T. Mattner, ISBN 978-0-9805142-1-6 (2008).

[19] M. L. Ekiel-Jeżewska and B. U. Felderhof, Periodic sedimentation of three particles in periodic boundary conditions, *Phys. Fluids* **17**, 093102 (2005).

[20] M. L. Ekiel-Jeżewska and B. U. Felderhof, Clusters of particles falling in a viscous fluid with periodic boundary conditions, *Phys. Fluids*, **18**, 121502 (2006).

[21] S. Jung, S. E. Spagnolie, K. Parikh, M. Schelley, and A.-K. Tornberg, Periodic sedimentation in a Stokesian fluid, *Phys. Rev. E* **74**, 035302(R) (2006).

[22] Cichocki B., Felderhof B. U., Hinsen K., Wajnryb E. and Bławzdziewicz J., J. Chem. Phys. **100** (1994), 3780.

[23] Ekiel-Jeżewska M. L. and Wajnryb E., *in: Theoretical Methods for Micro Scale Viscous Flows*, F. Feuillebois and A. Sellier, eds., Transworld Research Network, Kerala, 2009.

[24] B. Cichocki, M.L. Ekiel-Jeżewska and E. Wajnryb, J. Chem. Phys. **111**, 3265 (1999).